



Antioxidant and Anticancer Potential of Bioactive Compounds from Locally Isolated Microalgae

Ain Nafiza Basha¹, Fazrena Nadia Md Akhir^{1,*}, Nor'azizi Othman², Hirofumi Hara³

¹ Department of Chemical and Environmental Engineering, Malaysia-Japan International Institute of Technology, Universiti Teknologi Malaysia, Jalan Sultan Yahya Petra, 54100 Kuala Lumpur, Malaysia

² Department of Mechanical Precision Engineering, Malaysia-Japan International Institute of Technology, Universiti Teknologi Malaysia, Jalan Sultan Yahya Petra, 54100 Kuala Lumpur, Malaysia

³ Department of Biotechnology, Graduate School of Agricultural and Life Sciences, The University of Tokyo, 1-1-1, Yayoi, Bunkyo-ku, Tokyo, 113-8657, Japan

ARTICLE INFO

Article history:

Received 1 July 2024

Received in revised form 15 July 2024

Accepted 10 August 2024

Available online 16 August 2024

Keywords:

Microalgae; antioxidant; anticancer;
MCF-7; *Acutodesmus obliquus*;
Desmodesmus perforates

ABSTRACT

The current cancer therapy research focuses on the curative strategies but also targeting to improve the development of the preventive strategies. Anticancer, antibacterial, antioxidant, and anti-inflammatory effects have been demonstrated by the bioactive compound extracted from microalgae. Marine organisms, particularly microalgae might be one the alternative that may contribute to the achievement of one of Sustainable Development Goals (SDG) which is SDG 3, Good Health and Wellbeing. It is crucial for long-term sustainable development to ensure healthy lives and promote well-being at all ages. Two microalgal samples from Malaysia *Acutodesmus obliquus* (CN01) and *Desmodesmus perforates* (SP04) were isolated from fresh waters and cultivated. A total of three different solvents, polar, intermediate and nonpolar solvent was used to extracts the microalgae sample which are methanol, acetone and hexane, respectively. *Acutodesmus obliquus* and *Desmodesmus perforates* extracts show low total phenolic content in range from 0.2447 ± 0.0009 to 0.2614 ± 0.0003 . Secondly, the antioxidants activity of both microalgae shows significant radical scavenging activity especially in acetone extracts of both microalgae, with values of 11.5433 ± 0.0115 and 13.5847 ± 0.0618 for *Acutodesmus obliquus* (CN01) whereas *Desmodesmus* (SP04), 13.7898 ± 0.2601 and 22.3212 ± 0.2302 scavenging activity percentage. Then, the anticancer effect of the crude extracts was tested against breast cancer, MCF-7 breast cancer cell lines by using MTT assay and apoptosis assay. Methanolic extract has demonstrated high antiproliferative effect against MCF-7, with cell inhibition percentage within range of 87 ± 1.1271 % to 73 ± 0.2744 % and 82 ± 0.0236 % to 73 ± 0.0423 % of *Acutodesmus obliquus* (CN01) and *Desmodesmus* (SP04) respectively. Finally, both microalgae show similar compound group as HPLC is used to identified lipid, which is believed to play the big role in contributing to anticancer and antioxidant activity. This suggests that *Acutodesmus obliquus* sp and *Desmodesmus perforates* sp possess diverse biosynthetic capacities, potentially enabling the creation of new pharmaceuticals.

* Corresponding author.

E-mail address: fazrena@utm.my

<https://doi.org/10.37934/jhqol.3.1.4054>

1. Introduction

According to Globocan in 2020, cancer caused approximately 10 million deaths worldwide and 19,292,789 new cases were reported [1]. In Malaysia, cancer accounted for about 0.09% of the population's deaths, making it the leading cause of death in 2016, with 12.6% of deaths in government hospitals and 26.7% in private hospitals. The trend of cancer-related deaths in Malaysia increased from 11.3% in 2007 to 12.6% in 2016 [2].

Cancer occurs due to uncontrolled cellular proliferation [3], and current treatments using synthetic compounds are less efficient compared to natural compound-based drugs, which are easier to metabolize. Major treatments include alkylating agents and antimetabolites, which have significant side effects. Cancer can develop due to disruptions in cell death mechanisms, such as apoptosis, which is crucial for cancer therapy. The protein caspase plays a key role in apoptosis, and the tumor suppressor gene p53 helps prevent cancer. When p53 is mutated, its function is lost, leading to cancer formation [4].

Studies have shown that macroalgae and microalgae have potential anti-cancer properties, inducing apoptosis in cancer cells [5-7]. Microalgae, such as *Nannochloropsis*, contain various bioactive compounds like pigments, amino acids, and ω -3 fatty acids, which exhibit antitumor, antioxidant, and anti-inflammatory activities [8].

Up till today, cancer treatments conducted involve chemotherapeutic drugs, which cause severe side effects. Recent research focuses on natural ingredients from microorganisms, animals, and plants that target tumor cells without harming normal cells. Single-pathway treatments show limited success, necessitating combination therapies and advancements in precision medicine [9]. Nanomedicine can improve targeting but faces challenges like high interstitial fluid pressure and drug delivery issues in certain cancers [10]. Therefore, the extraction and analysis of bioactive compounds from natural sources should be considered as new alternative [11,12]. Plants and algae offer promising sources of anticancer agents with fewer side effects as compare to today treatment such as chemotherapy [13]–[15]. Algae, particularly microalgae, show potential in inducing cancer cell apoptosis without affecting normal cells, suggesting a need for more research into their use in cancer treatment [16]. For example, *Euglena tuba* extract has been effective against cancer cells while sparing normal cells [17].

There are over 200 types of cancer, which can spread to other tissues through metastasis, often resulting in death. Cancer develops due to damaged DNA, faulty DNA repair mechanisms, activation of tumors, and loss of tumor suppression functions [18]. Major causes of cancer include an unhealthy lifestyle, poor diet, aging, smoking, and environmental pollutants. High fatality rates often result from therapy failure and tumor relapse. During treatment, cancer cells may develop resistance due to improper pharmacological strategies or various drug resistance mechanisms such as drug efflux, detoxification, changes in drug targets, effective DNA repair, blocked apoptosis, and the development of resistant cancer stem cells [19].

Microalgae have significant potential for human health and medicine, with applications in cardiovascular health, cancer treatment, inflammation reduction, infection prevention, immune enhancement, and cholesterol reduction. They are also useful in treating tumors, stomach ulcers, and aiding wound healing, with components capable of detoxifying the body [20,21].

2. Methodology

2.1 Microalgae Cultivation

Microalgae culture for this study was obtained from an isolated strain sourced from the Algal Biomass Research Lab, *Acutodesmus obliquus* (CN01, MJIIT-UTM, Malaysia) and *Desmodesmus* (SP04, MJIIT-UTM). Since microalgae require specific nutrients for healthy growth, AF6 and BG-11 media were prepared for *Acutodesmus obliquus* sp and *Desmodesmus* sp respectively. Subsequently, the microalgae were allowed to grow under optimal conditions, exposed to white, fluorescent light at 80 μ mol photons/m²/s for a 12-hour photoperiod at 25°C, with continuous aeration. Regular monitoring of the microalgal culture for growth and changes in cell density was conducted using a UV-VIS spectrophotometer (Hach DR 6000, Hach, U.S.A.) at 750 nm optical density (OD). Harvesting of the microalgae occurred during the log phase, whereby harvested microalgae were stored at -80°C until further procedures were conducted. Subsequently, the frozen microalgae underwent freeze-drying using a freeze dryer (Scientz-10ND) to remove excess water. The resulting freeze-dried microalgae were stored at -80°C to preserve their viability until further use [23,24].

2.2 Solvent Extraction

5 grams of freeze-dried *A. obliquus* or *Desmodesmos* powder were agitated in 100 mL of hexane, acetone, or methanol for 24 hours, followed by filtration through cotton. The solvent was subsequently evaporated using a spin dryer at 35°C, and the resulting dried extract was stored at -20°C for future applications [25-27].

2.3 Measurement of Phenolic Content

The measurement of phenolic content of microalgae extract is done by using colorimetric assay. In this study, a suitable phenolic standard stock solution is prepared, with gallic acid serving as the chosen standard for analysis. Subsequently, a reaction mixture comprising crude extracts, Folin-Ciocalteu reagent, and sodium carbonate solution is meticulously prepared. Following preparation, the mixture is allowed to incubate for a duration of 30 minutes to facilitate the reaction process. After the incubation period, the absorbance of the resulting mixture is measured at 750nm using spectrometer UV-VIS spectrophotometer (Hach DR6000). Finally, the phenolic content of the microalgae extract is determined by calculating it based on the absorbance values obtained from the analysis, enabling a comprehensive assessment of the extract's phenolic composition [22].

2.4 Radical Scavenging Inhibition Activity

In this study, DPPH (2,2-diphenyl-1-picrylhydrazyl) assay method is used to evaluate the antioxidant content of microalgae. To initiate the antioxidant activity assessment, a 1mM DPPH solution is first prepared by dissolving 1.2mg of DPPH in 50ml of methanol within a volumetric flask. The absorbance of this solution is meticulously evaluated and adjusted to fall within the prescribed range of 0.65 $\pm$ 0.02 at 517nm. Following this, compounds extracted from the microalgae are diluted to the desired concentration range of 2mg/ml to 10mg/ml using methanol as the selected solvent. Subsequently, 0.8ml of the prepared DPPH solution is mixed with 0.2ml of the diluted microalgae extract in a 1ml centrifuge tube, and the resulting mixture is then incubated for a duration of 30 minutes at room temperature under dark conditions. After the incubation period, the absorbance of the reaction mixture is measured at a wavelength of 517nm by using UV-VIS spectrophotometer

(Hach DR6000). Same procedure is repeated from ascorbic acid standard. This method allows for the comprehensive assessment of the microalgae extract's radical scavenging potential, providing valuable insights into its antioxidant properties [28]. Utilizing the acquired absorbance values, the antioxidant activity of the microalgae extract is determined by applying the formula below Equation (1), which calculates the radical scavenging activity (RSA) as a percentage: where Abs_{DPPH} represents Absorbance value of standard DPPH solution, and Abs_{sample} represents Absorbance value of standard DPPH solution.

$$RSA \% = \left(\frac{Abs_{DPPH} - Abs_{sample}}{Abs_{DPPH}} \right) \times 100 \quad (1)$$

2.5 Cell Culture

MCF-7 is a breast cancer cell line that is used in vitro in this study. Sterile techniques and proper safety precautions were involved throughout the entire process to prevent any contamination as referred to [29]. The cell line MCF-7 is cultured using Minimum Essential Medium (MEM) supplemented with 10% fetal bovine serum (FBS), 1% L-glutamine, and 1% penicillin/streptomycin to support cell growth and proliferation. The cells are then incubated in a humidified incubator with 5% CO₂ at 37°C by using Astec, APC-50DR. Daily monitoring of cell growth is conducted using an inverted microscope (Olympus CKX41), and the medium is changed every 3-5 days or as needed to maintain optimal growth conditions.

2.6 MTT Assay

MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay is used in this study in which colorimetric assay is used to measure cell viability and proliferation in vitro. The experimental procedure begins by plating cells in a 96-well plate at a density of 1×10^5 cells per well. Following plating, the cells are incubated at 37°C in a humidified incubator with 5% CO₂ for approximately 48 to 72 hours until firm attachment to the plate is confirmed. Upon confirmation of cell attachment, the medium is aspirated from the wells and replaced with microalgae extract solutions of varying concentrations, while fresh medium serves as a negative control and DMSO acts as a positive control. Subsequently, the cells are further incubated for an additional 24 hours. Following this incubation period, the MTT solution is prepared by dissolving MTT in phosphate-buffered saline (PBS) at a concentration of 5 mg/ml. After the 24-hour incubation period, the extracts, medium, and DMSO are removed from the wells, and an equal volume of MTT solution is added. The cells are then incubated for 4 hours at 37°C in a humidified incubator with 5% CO₂. Following the incubation, the MTT solution is carefully aspirated, and DMSO is added to dissolve the formazan crystals formed by viable cells. Finally, the absorbance of the dissolved formazan is measured at 560 nm wavelengths using a microplate reader, GloMax® Discover (PromegaInstrument, 9700000313) allowing for the assessment of cell viability and potential effects of the microalgae extract [30]. The cell viability and proliferation of the cells is calculated using the absorbance values obtained and formula as shown below Equation (2). Where OD_{sample} represents Optical density of sample, and $OD_{control}$ represents Optical density of control.

$$Cell\ viability\ percentage\ (\%) = \frac{OD_{sample}}{OD_{control}} \times 100 \quad (2)$$

2.7 Phytochemical Identification using GCMS

The chemical components of the extracts from both microalgae were analyzed using GC-MS (The Agilent 7820A GC) featuring a fused silica capillary column (30 m × 0.25 mm × 0.25 µm, coated with DB-5) and an electron ionization (EI) operating at 70 eV. The injector temperature was maintained at 250°C. The oven temperature was programmed to start at 40°C for 1 minute and then increase at a rate of 3°C per minute until reaching 280°C. Helium served as the carrier gas at a flow rate of 1 mL/min. Identification of compounds was achieved by comparing their mass spectra with those in the National Institute of Standards and Technology (NIST) library [31].

2.8 Statistical Analysis

The experiments were conducted three or more times, and the results are presented as mean ± SD. Group differences were assessed using one-way ANOVA with GraphPad Prism 9.0 software, where significance levels were denoted as ****P < .0001.

3. Result and Discussion

3.1 Total Phenolic Content

Phenolic compounds are key antioxidants, stabilizing radicals by donating hydrogen atoms or electrons. The total phenolic content in microalgae was measured as gallic acid equivalent (GAE) to compare different strains. These compounds, significant for their scavenging potential due to hydroxyl groups, help microalgae mitigate oxidative stress [32-34]. Therefore, in this study two local microalgae are used to analyse the total phenolic content which is a crucial parameter for measuring the antioxidant potential of *Acutodesmus obliquus* CN01 and *Desmodesmus* SP04. Total phenolic compound is determined by using Folin-Ciocalteu method. Total phenolic compounds are measured using the Folin-Ciocalteu method, where microalgae extract react with Folin-Ciocalteu reagent to form a blue complex. The phenolic content is determined by measuring the colour intensity with a spectrophotometer, using a gallic acid standard curve.

Three different solvents with different polarity were used to evaluate the total phenolic content (TPC) which are methanol, acetone and hexane of two microalgae which are *Acutodesmus obliquus* (CN01) and *Desmodesmus* (SP04). From Figure 1, *Acutodesmus obliquus* (CN01) has a total phenolic content of 0.2447 to 0.2522 mg/g GAE, while *Desmodesmus* (SP04) ranges from 0.2466 to 0.2614 mg/g GAE. The highest phenolic content is found in *Desmodesmus* methanolic extract (0.2614 ± 0.003 mg/g GAE), followed by *Acutodesmus obliquus* methanolic extract (0.2522 ± 0.0007 mg/g GAE). Hexane extracts show the lowest TPC values for both microalgae. Acetone extracts have higher TPC in *Desmodesmus*, 0.2487 ± 0.0001 mg/g GAE compared to *Acutodesmus obliquus*, 0.2472 ± 0.0003 mg/g GAE. Overall, TPC is highest in methanol extracts, followed by acetone, and lowest in hexane. This trend aligns with the polar nature of phenolic compounds, as seen in previous studies [32,35].

Comparing our analyzed strain of *Acutodesmus obliquus* (CN01) and *Desmodesmus* (SP04), phenolic content is significantly lower compared to other previously published data. For example, green microalga *Scenedesmus* sp was observed to have the highest TPC in ethanol/water, ranging from 0.5 to 4.6 mg GAE g⁻¹ compared to TPC to 0.2447 mg/g to 0.2522 mg GAE g⁻¹ in *Scenedesmus* sp. It's crucial to acknowledge that various factors such as growth conditions, pH, temperature, light exposure, harvesting timing, alongside inherent variability and extraction techniques, influence the phenolic content and antioxidant capacities of microalgae [36]. A lot of factors play such a big role in extracting phenolics compound, as shown in Ramalingam Parthasarathy study, higher phenolic

compound of 3.4 mg/gae can be extracted when using hot water extraction method. Thus, explained that different methods and solvents do affect the phenolic content extracted.




Fig. 1. Total Phenolic Content of *A. obliquus* and *Desmodesmus* in methanol, acetone and hexane. Expressed as mean $\pm$ standard deviation (SD) of triplicate sample extract

3.2 Radical Scavenging Activity

The DPPH assay evaluates the ability of tested antioxidants to reduce the DPPH $\cdot$ radical either through direct electron transfer or by quenching the radical via hydrogen atom transfer [37]. The solubility of antioxidants, primarily influenced by the polarity of the solvent employed, plays a crucial role in accurately evaluating antioxidant capacity. In this research, we utilized three solvents with differing polarities: methanol, acetone, and hexane for extraction.

The antioxidant potential of *Acutodesmus obliquus* (CN01) and *Desmodesmus* (SP04), extracts was assessed through DPPH assays. Figure 2 and Figure 3 presents the findings, indicating the percentage of DPPH radical scavenging activity of microalgae. The results reveal that the highest antioxidant capacity, was observed in the acetone, with values of 11.5433 ± 0.0115 and 13.5847 ± 0.0618 for *Acutodesmus obliquus* (CN01) whereas *Desmodesmus* (SP04), 13.7898 ± 0.2601 and 22.3212 ± 0.2302 scavenging activity percentage. Extracts in other solvents demonstrated comparatively lower antioxidant capacities. Statistical analysis showed significant differences ($p < 0.0001$) between hexane, methanol, and acetone compared to ascorbic acid. Our findings align with prior research indicating that the selection of solvent significantly impacts the extraction yield of substances, as suggested by [35].

Between *Acutodesmus obliquus* (CN01) and *Desmodesmus* (SP04), *Desmodesmus* (SP04), shown high significance scavenging activity compared to *Acutodesmus obliquus* (CN01) in overall as shown in Figure 2 and Figure 3. However, both microalgae do display the same pattern of which acetone exhibit higher scavenging activity than both methanol and hexane. conversely, the ethanolic extracts from *U. fasciata* and *G. salicornia* showed significant antioxidant activity. The antioxidant effects can be credited to phenolic compounds along with various other phytochemical compounds like carotenoids, polyunsaturated fatty acids (PUFA), and polysaccharides found within the algal extract, [22].

Martin Almendinger *et al.*, [38] discovered that the strains with the highest content of water-soluble compounds were measured between 12–14 μ mol ascorbic acid equivalent per gram, while

N. oleobundans and *A. obliquus* were the strains exhibiting the highest concentrations of lipid-soluble compounds, ranging between 60–80 μmol trolox equivalent per gram. According to Amzah *et al.*, [39], this particular strain of *Acutodesmus obliquus*, also referred to as *Scenedesmus obliquus* or *Tetradesmus obliquus*, has been previously demonstrated to possess a significant number of fatty acids. Additional elements like the composition of fatty acids, could potentially enhance the antioxidant capacity of both *Acutodesmus obliquus* (CN01) and *Desmodesmus* (SP04), as indicated [40,41].




Fig. 2. DPPH radical scavenging activity of *Acutodesmus obliquus* of methanol, acetone, and hexane extracts. All result is expressed in mean $\pm$ SD. ($p < 0.0001$) ($n = 3$)




Fig. 3. DPPH radical scavenging activity of *Desmodesmus* of methanol, acetone, and hexane extracts. All result is expressed in mean $\pm$ SD. ($p < 0.0001$) ($n = 3$)

3.3 Anticancer Property

The MTT assay is used to evaluate cytotoxicity in MCF-7 cell cultures. It involves adding the MTT compound to cells, which is metabolized by living cells into insoluble formazan crystals. After a period of incubation, the cells are lysed to release the formazan, which is then solubilized and measured spectrophotometrically. The absorbance is directly proportional to the number of viable cells, allowing for the assessment of cytotoxic effects of substances or treatments on cell cultures [28,42].

The cytotoxicity impact of three solvent namely methanol, acetone and hexane extract of *Acutodesmus obliquus* (CN01) and *Desmodesmus* (SP04) was examined on MCF-7 breast cancer cell line via the colorimetric MTT assay. Where DMSO is use a control standard to compare with the extracts. Both microalgae show same patter in which MCF-7 cell viability decreases as concentration of the extract increases, in other word, all extract shows the same pattern of dose dependent manner against MCF-7.

Based on Figure 4 and Figure 5, acetone extracts from both microalgae showed the lowest cytotoxicity on MCF-7 cells: $83 \pm 1.0723\%$ to $79 \pm 0.6934\%$ for *Acutodesmus obliquus* (CN01) and $81 \pm 0.0473\%$ to $76 \pm 0.5271\%$ for *Desmodesmus* (SP04). Methanol extracts recorded the highest cell inhibition: $87 \pm 1.1271\%$ to $73 \pm 0.2744\%$ for *Acutodesmus obliquus* (CN01) and $82 \pm 0.0236\%$ to $73 \pm 0.0423\%$ for *Desmodesmus* (SP04). *Desmodesmus* had a higher anti-proliferative effect on MCF-7 cells than *Acutodesmus obliquus* (CN01), especially in methanolic extracts. The anticancer effect of

the extracts follows the order: methanol > acetone > hexane. These findings align with studies by Elia et al. [43] showing acetone extracts of *Laurencia papillosa* having dose-dependent cytotoxic effects on MCF-7 cells, and by Goh et al., [44] and Suh et al., [45], demonstrating the anticancer properties of ethyl acetate extracts from various algae.

Whereas methanolic extracts of both microalgae shows significant anticancer effect on MCF-7 demonstrates the decrease in cell viability by 73 ± 0.2744 % and 73 ± 0.0423 % of 8 mg/ml *Acutodesmus obliquus* (CN01) and *Desmodesmus* (SP04) respectively. Similar observations have been made in the methanol extracts obtained from *Dunaliella tertiolecta* were effective solely against the A2058 human melanoma cell line. Moreover, Nigjeh et al., [46] shows that the impact of crude ethanol extracts derived from a native microalga, *Chaetoceros calcitrans*, on human breast cell lines (MCF-7) demonstrate significant cytotoxic properties and might play the role in inducing the mechanism of apoptosis. Differences in the cytotoxic impact of a particular algal species can be attributed to the selection of extraction solvent and its efficacy in extracting bioactive compounds, bioactive compound composition, alongside seasonal and geographical fluctuations influenced by environmental factors like temperature, light exposure, and salinity alterations [47,48].




Fig. 4. Cytotoxicity of *Acutodesmus obliquus* (CN01) in human breast cell line MCF-7 cells, following incubation for 24 hours, using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. All result is expressed in mean $\pm$ SD. ($p < 0.0001$) ($n = 3$)




Fig. 5. Cytotoxicity of *Desmodesmus* (CN01) in human breast cell line MCF-7 cells, following incubation for 24 hours, using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. All result is expressed in mean $\pm$ SD. ($p < 0.0001$) ($n = 3$)

3.4 Phytochemical Analysis by GCMS

Two extracts out of six extracts have been chosen based on the previous antioxidant capacity and anticancer property, both methanolic extract of *Acutodesmus obliquus* (CN01) and *Desmodesmus* showing the most effective result in reducing the cell proliferation of MCF-7 breast cancer cell line. Therefore, further investigation of the phytochemical constituents is carried out using GCMS. As mentioned in the literature found, generally, green algae contain alkaloids, phenols, flavonoids, lipids, and polysaccharides which demonstrate anti-cancer properties and offer health benefits associated with antioxidants.

From GCMS result, we found that both freshwater green microalgae's major content is lipid. Fatty acids represent a significant lipid category present in macroalgae, serving as essential components of cell membranes such as phosphoglycerides or as reserves for energy for instance, triacylglycerol. Typically, they exhibit higher concentrations compared to other lipids such as sterols, hydrocarbons, and fatty alcohols. When comparing our findings with existing literature, it was noted that the predominant fatty acids detected in *Acutodesmus obliquus* (CN01) and *Desmodesmus* (SP04) which are fatty acid, fatty acid methyl ester, fatty alcohol were consistent with those previously reported [49].

As referring to the finding tabulated in Table 1 and Table 2, the phytochemical found in the microalgae extracted, both *Acutodesmus obliquus* (CN01) and *Desmodesmus* (SP04) was compared with previous studies. This shown that Dodecane and tetradecane from *Aspergillus niger* AK6 exhibit anticancer properties by inducing apoptosis and necrosis in MCF-7 cells, and also have antimicrobial and antifungal effects [50]. Dodecene from *X. testudinaria* shows anticarcinogenic effects on HeLa cells [51]. Metabolites like 1-Nonadecene, 1-Octadecene, and 1-Tetradecene in algae and plants combat cancer, act as antioxidants, and have antimicrobial properties [52]. Tetradecane in macroalgae contributes to antioxidant and antimicrobial effects [53]. *S. platensis* has tetradecane as a major component with antimicrobial activity [54]. Tetradecane from *Cladostephus spongiosus*

shows antioxidant properties [55]. Cyclotetradecane demonstrates antimicrobial [56] and antioxidant potential [57]. Tetradecane from *Ulothrix* sp shows antifungal activity.

Pacheco *et al.*, [58] reported that fatty acid extract of *A. utricularis* has antiproliferative activity on MCF-7 and MDA-MB-231 breast cancer cells. Hexadecanoic acid, or palmitic acid, enhances cytotoxic effects against cancer cells [59]. Octadecanol has antimicrobial and antioxidant properties [60].

GC-MS analysis of *Acutodesmus obliquus* (CN01) extract revealed bioactive compounds, including hexadecenoic acid, heptadecanoic acid, and octadecanoic acid. Lipid from *Chlorella sorokiniana* shows high cytotoxicity to A375, A549, and HeLa cells [61]. Hexadecanoic acid from *T. ornate* inhibits colon cancer cell proliferation [62]. Fatty acids from *C. silvatica* methanol extract reduce HeLa, PC3, and MCF7 cancer cell viability. Freshwater microalgae lipids have antimicrobial properties by damaging cell membranes [63].

9,12,15-Octadecatrienoic acid, methyl ester, from *Acutodesmus obliquus* (CN01), known as alpha-linolenic acid (ALA), is a precursor for n-3 polyunsaturated fatty acids (PUFAs) [64]. ALA exhibits potent antiproliferative effects on MCF-7 cells [65], reducing neoplastic transformation, inhibiting growth, and inducing apoptosis. ALA decreases cell viability and alters gene expression in various breast cancer cells [66]. ALA suppresses estrogen receptor-positive MCF-7 cell growth both in vitro and in vivo, likely by modulating ER-related signaling pathways [67]. These PUFAs may significantly contribute to the anticancer effect on MCF-7 cells in this research.

Table 1

Compounds found in methanolic extract of *Acutodesmus obliquus* (CN01) - Name, Retention Time (RT), and Peak area percentage of compounds present in the sample during GCMS analysis

Compound	Retention time (min)	Area %	Molecular weight (g/mol)	Molecular formula
7,10,13-Hexadecatrienoic acid, methyl ester	24.557	4.66	264.4	C ₁₇ H ₂₈ O ₂
Hexadecanoic acid, methyl ester	25.046	11.28	270.5	C ₁₇ H ₃₄ O ₂
Heptadecanoic acid, methyl ester	26.777	4.74	284.5	C ₁₈ H ₃₆ O ₂
9,12-Octadecadienoic acid (Z,Z)-, methyl ester	28.020	4.46	294.5	C ₁₉ H ₃₄ O ₂
9,12,15-Octadecatrienoic acid, methyl ester	28.223	15.19	292.5	C ₁₉ H ₃₂ O ₂
Octadecanoic acid, methyl ester	28.556	4.01	298.5	C ₁₉ H ₃₈ O ₂

Table 2

Compounds found in methanolic extract of *Desmodesmus* (SP04) - Name, Retention Time (RT), and Peak area percentage of compounds present in the sample during GCMS analysis

Compound	Retention time (min)	Area %	Molecular weight (g/mol)	Molecular formula
2-Dodecene	16.861	2.57	168.32	C ₁₂ H ₂₄
Dodecane	21.944	5.03	170.33	C ₁₂ H ₂₆
1-Tridecene	26.136	9.58	182.35	C ₁₃ H ₂₆
Cyclododecane	30.496	8.48	168.32	C ₁₂ H ₂₄
Tetradecane	30.825	4.16	198.39	C ₁₄ H ₃₀
1-Pentadecene	34.639	6.77	210.40	C ₁₅ H ₃₀
Pentadecane	34.947	3.15	212.41	C ₁₅ H ₃₂
1-Octadecanol	42.310	1.96	270.5	C ₁₈ H ₃₈ O

4. Conclusions

The methanolic extracts of *Acutodesmus obliquus* (CN01) and *Desmodesmus* (SP04) demonstrated significant anticancer properties by effectively reducing the proliferation of MCF-7 breast cancer cells. GC-MS analysis revealed that these extracts are rich in bioactive lipids, including fatty acids, fatty acid methyl esters, and fatty alcohols, which are known for their anticancer, antioxidant, and antimicrobial properties. These findings align with existing literature, underscoring the therapeutic potential of green microalgae-derived compounds. This study indicated that these lipids play a crucial role in inducing apoptosis and inhibiting cancer cell proliferation through various cellular pathways. The identified compounds, such as dodecane, tetradecane, and alpha-linolenic acid (ALA), exhibited potent antiproliferative effects and contributed to the overall anticancer activity of the extracts. These bioactive lipids not only inhibit cancer growth but also modulate gene expression and cellular signalling pathways, enhancing their therapeutic efficacy. Future research should focus on comprehensive phytochemical profiling, in vivo validation, and comparative analysis with other anticancer agents. Additionally, exploring other microalgae species and developing advanced formulations can further enhance the stability, bioavailability, and targeted delivery of these bioactive compounds. By pursuing these avenues, we can unlock the full potential of green microalgae-derived compounds in cancer therapy and develop novel, effective anticancer agents.

Acknowledgement

This research was funded by Malaysia Japan International Institute of Technology Incentive.

References

- [1] Sung, Hyuna, Jacques Ferlay, Rebecca L. Siegel, Mathieu Laversanne, Isabelle Soerjomataram, Ahmedin Jemal, and Freddie Bray. "Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries." *CA: a cancer journal for clinicians* 71, no. 3 (2021): 209-249. <https://doi.org/10.3322/caac.21660>
- [2] Azizah, A. M., A. Noor Hashimah, and K. M. Yusof. "Malaysian study on cancer survival (myscan)." *National Cancer Institute* 4 (2018): 10-20.
- [3] Offer, Steven M., and Robert B. Diasio. "Pharmacogenomics and the role of genomics in cancer therapeutics." *Cancer Chemotherapy, Immunotherapy and Biotherapy: Principles and Practice* (2018).
- [4] Taherzadeh-Soureshjani, Parisa, and Mohammad Chehelgerdi. "Algae-mediated route to cuprous oxide (Cu₂O) nanoparticle: differential expression profile of MALAT1 and GAS5 lncRNAs and cytotoxic effect in human breast cancer." *Cancer Nanotechnology* 11 (2020): 1-34. <https://doi.org/10.1186/s12645-020-00066-4>
- [5] Sit, Nam Weng, Yik Sin Chan, S. C. Lai, Lay Ngor Lim, G. T. Looi, P. L. Tay, Y. T. Tee, Y. Y. Woon, Kong Soo Khoo, and Hean Chooi Ong. "In vitro antidermatophytic activity and cytotoxicity of extracts derived from medicinal plants and marine algae." *Journal de Mycologie Medicale* 28, no. 3 (2018): 561-567. <https://doi.org/10.1016/j.mycmed.2018.07.001>
- [6] Senousy, Hoda H., Sawsan Abd Ellatif, and Shafaqat Ali. "Assessment of the antioxidant and anticancer potential of different isolated strains of cyanobacteria and microalgae from soil and agriculture drain water." *Environmental Science and Pollution Research* 27 (2020): 18463-18474. <https://doi.org/10.1007/s11356-020-08332-z>
- [7] Goncu, Beyza, Ece Sevgi, Cagla Kizilarlan Hancer, Guzin Gokay, and Nur Ozten. "Differential anti-proliferative and apoptotic effects of lichen species on human prostate carcinoma cells." *PLoS One* 15, no. 9 (2020): e0238303. <https://doi.org/10.1371/journal.pone.0238303>
- [8] Pangestuti, Ratih, and Se-Kwon Kim. "Biological activities and health benefit effects of natural pigments derived from marine algae." *Journal of functional foods* 3, no. 4 (2011): 255-266. <https://doi.org/10.1016/j.jff.2011.07.001>
- [9] Zorofchian Moghadamtousi, Soheil, Hamed Karimian, Ramin Khanabdali, Mahboubeh Razavi, Mohammad Firoozinia, Keivan Zandi, and Habsah Abdul Kadir. "Anticancer and antitumor potential of fucoidan and fucoxanthin, two main metabolites isolated from brown algae." *The Scientific World Journal* 2014, no. 1 (2014): 768323. <https://doi.org/10.1155/2014/768323>
- [10] Zugazagoitia, Jon, Cristiano Guedes, Santiago Ponce, Irene Ferrer, Sonia Molina-Pinelo, and Luis Paz-Ares. "Current challenges in cancer treatment." *Clinical therapeutics* 38, no. 7 (2016): 1551-1566.

- <https://doi.org/10.1016/j.clinthera.2016.03.026>
- [11] Ismail, Husna Filzah, Fazrena Nadia Md Akhir, and Hirofumi Hara. "Optimization of Extraction Solvents on the Antioxidant Properties of Coconut Waste." *Journal of Advanced Research in Fluid Mechanics and Thermal Sciences* 117, no. 1 (2024): 109-117. <https://doi.org/10.37934/arfmts.117.1.109117>
- [12] Basha, Ain Nafiza, Fazrena Nadia Md Akhir, and Hirofumi Hara. "Anticancer Potential of Bioactive Compounds from Microalgae. A Review." *Journal of Advanced Research in Micro and Nano Engineering* 20, no. 1 (2024): 1-9. <https://doi.org/10.37934/armne.20.1.19>
- [13] Gaumet, Marie, Angelica Vargas, Robert Gurny, and Florence Delie. "Nanoparticles for drug delivery: the need for precision in reporting particle size parameters." *European journal of pharmaceuticals and biopharmaceutics* 69, no. 1 (2008): 1-9. <https://doi.org/10.1016/j.ejpb.2007.08.001>
- [14] Provenzano, Paolo P., Carlos Cuevas, Amy E. Chang, Vikas K. Goel, Daniel D. Von Hoff, and Sunil R. Hingorani. "Enzymatic targeting of the stroma ablates physical barriers to treatment of pancreatic ductal adenocarcinoma." *Cancer cell* 21, no. 3 (2012): 418-429. <https://doi.org/10.1016/j.ccr.2012.01.007>
- [15] Jain, Rakesh K., and Triantafyllos Stylianopoulos. "Delivering nanomedicine to solid tumors." *Nature reviews Clinical oncology* 7, no. 11 (2010): 653-664. <https://doi.org/10.1038/nrclinonc.2010.139>
- [16] Abd El-Hack, Mohamed E., Sameh Abdelnour, Mahmoud Alagawany, Mohamed Abdo, Moustafa A. Sakr, Asmaa F. Khafaga, Samir A. Mahgoub, Shaaban S. Elnesr, and Manar G. Gebriel. "Microalgae in modern cancer therapy: Current knowledge." *Biomedicine & pharmacotherapy* 111 (2019): 42-50. <https://doi.org/10.1016/j.biopha.2018.12.069>
- [17] Panja, Sourav, Nikhil Baban Ghate, and Nripenderanath Mandal. "A microalga, *Euglena tuba* induces apoptosis and suppresses metastasis in human lung and breast carcinoma cells through ROS-mediated regulation of MAPKs." *Cancer Cell International* 16 (2016): 1-13. <https://doi.org/10.1186/s12935-016-0330-5>
- [18] Vineis, Paolo, and Christopher P. Wild. "Global cancer patterns: causes and prevention." *The Lancet* 383, no. 9916 (2014): 549-557. [https://doi.org/10.1016/S0140-6736\(13\)62224-2](https://doi.org/10.1016/S0140-6736(13)62224-2)
- [19] Alves, Celso, Eurico Serrano, Joana Silva, Carlos Rodrigues, Susete Pinteus, Helena Gaspar, Luis M. Botana, Maria C. Alpoim, and Rui Pedrosa. "Sphaerococcus coronopifolius bromoterpenes as potential cancer stem cell-targeting agents." *Biomedicine & Pharmacotherapy* 128 (2020): 110275. <https://doi.org/10.1016/j.biopha.2020.110275>
- [20] de Jesus Raposo, Maria Filomena, Rui Manuel Santos Costa de Moraes, and Alcina Maria Miranda Bernardo de Moraes. "Health applications of bioactive compounds from marine microalgae." *Life sciences* 93, no. 15 (2013): 479-486. <https://doi.org/10.1016/j.lfs.2013.08.002>
- [21] Levine, Ira A. "Algae: A way of life and health." In *Microalgae in health and disease prevention*, pp. 1-10. Academic Press, 2018. <https://doi.org/10.1016/B978-0-12-811405-6.00001-3>
- [22] Arguelles, E. D. L. R., Antonio C. Laurena, Milagrosa R. Martinez-Goss, and Rosario G. Monsalud. "Antibacterial activity, total phenolic content and antioxidant capacity of a green microalga *Desmodesmus* sp.(U-AU2) from Los Baños, Laguna (Philippines)." *Journal of Nature Studies* 16, no. 2 (2017): 1-13.
- [23] Helamieh, Mark, Annkathrin Gebhardt, Marco Reich, Friederike Kuhn, Martin Kerner, and Klaus Kümmerer. "Growth and fatty acid composition of *Acutodesmus obliquus* under different light spectra and temperatures." *Lipids* 56, no. 5 (2021): 485-498. <https://doi.org/10.1002/lipd.12316>
- [24] Othman, Fatin Syahirah, Haryati Jamaluddin, Zaharah Ibrahim, Hirofumi Hara, Nurul Ashyikin Yahya, Koji Iwamoto, and Shaza Eva Mohamad. "Production of α -linolenic Acid by an Oleaginous Green Algae *Acutodesmus obliquus* Isolated from Malaysia." *Journal of Pure & Applied Microbiology* 13, no. 3 (2019). <https://doi.org/10.22207/JPAM.13.3.01>
- [25] Arceusz, Agnieszka, Marek Wesolowski, and Pawel Konieczynski. "Methods for extraction and determination of phenolic acids in medicinal plants: a review." *Natural product communications* 8, no. 12 (2013): 1934578X1300801238. <https://doi.org/10.1177/1934578X1300801238>
- [26] Ventura, S. P. M., B. P. Nobre, F. Ertekin, M. Hayes, M. García-Vaquero, F. Vieira, M. Koc, L. Gouveia, M. R. Aires-Barros, and A. M. F. Palavra. "Extraction of value-added compounds from microalgae." In *Microalgae-based biofuels and bioproducts*, pp. 461-483. Woodhead Publishing, 2017. <https://doi.org/10.1016/B978-0-08-101023-5.00019-4>
- [27] Gallego, Rocío, Lidia Montero, Alejandro Cifuentes, Elena Ibáñez, and Miguel Herrero. "Green extraction of bioactive compounds from microalgae." *Journal of Analysis and Testing* 2, no. 2 (2018): 109-123. <https://doi.org/10.1007/s41664-018-0061-9>
- [28] Haq, Samina Hyder, Ghaida Al-Ruwaished, Moudhi Abdullah Al-Mutlaq, Sundus Ali Naji, Maha Al-Mogren, Sarah Al-Rashed, Qura Tul Ain, Abir Abdullah Al-Amro, and Adnan Al-Mussallam. "Antioxidant, anticancer activity and phytochemical analysis of green algae, *Chaetomorpha* collected from the Arabian Gulf." *Scientific reports* 9, no. 1 (2019): 18906. <https://doi.org/10.1038/s41598-019-55309-1>
- [29] Mohamadnia, Sonia, Omid Tavakoli, Mohammad Ali Faramarzi, and Zahra Shamsollahi. "Production of fucoxanthin

- by the microalga *Tisochrysis lutea*: A review of recent developments." *Aquaculture* 516 (2020): 734637. <https://doi.org/10.1016/j.aquaculture.2019.734637>
- [30] Vasudevan, S., M. P. Arulmoorthy, and R. Suresh. "Isolation, purification and structural elucidation of secondary metabolites from *Microcystis aeruginosa* bloom from Muttukadu estuary and its in vitro antibacterial, antioxidant and anticancer potency." *South African journal of botany* 132 (2020): 59-67. <https://doi.org/10.1016/j.sajb.2020.03.030>
- [31] Olasehinde, Tosin A., Ejowwokoghene C. Odjadjare, Leonard V. Mabinya, Ademola O. Olaniran, and Anthony I. Okoh. "Chlorella sorokiniana and Chlorella minutissima exhibit antioxidant potentials, inhibit cholinesterases and modulate disaggregation of β -amyloid fibrils." *Electronic Journal of Biotechnology* 40 (2019): 1-9. <https://doi.org/10.1016/j.ejbt.2019.03.008>
- [32] Hajimahmoodi, Mannan, Mohammad Ali Faramarzi, Najmeh Mohammadi, Neda Soltani, Mohammad Reza Oveisi, and Nastaran Nafissi-Varcheh. "Evaluation of antioxidant properties and total phenolic contents of some strains of microalgae." *Journal of Applied Phycology* 22 (2010): 43-50. <https://doi.org/10.1007/s10811-009-9424-y>
- [33] Safaar, Hamed, Jonathan Van Wagenen, Per Møller, and Charlotte Jacobsen. "Carotenoids, phenolic compounds and tocopherols contribute to the antioxidative properties of some microalgae species grown on industrial wastewater." *Marine drugs* 13, no. 12 (2015): 7339-7356. <https://doi.org/10.3390/md13127069>
- [34] Taş, Beyhan, Ömer Ertürk, Özlem Yılmaz, Melek Çol Ayvaz, and Emine Yurdakul Ertürk. "Chemical components and biological activities of two freshwater green algae from Ordu, Turkey/Türkiye'nin Ordu ilinden iki tatlı su yeşil alglerinin kimyasal bileşenleri ve biyolojik aktiviteleri." *Turkish Journal of Biochemistry* 40, no. 6 (2015): 508-517. <https://doi.org/10.1515/tjb-2015-0042>
- [35] Jerez-Martel, Idaira, Sara García-Poza, Gara Rodríguez-Martel, Milagros Rico, Cristina Afonso-Olivares, and Juan Luis Gómez-Pinchetti. "Phenolic profile and antioxidant activity of crude extracts from microalgae and cyanobacteria strains." *Journal of Food Quality* 2017, no. 1 (2017): 2924508. <https://doi.org/10.1155/2017/2924508>
- [36] Bulut, Onur, Dilan Akin, Çağla Sönmez, Ayşegül Öktem, Meral Yücel, and Hüseyin Avni Öktem. "Phenolic compounds, carotenoids, and antioxidant capacities of a thermo-tolerant *Scenedesmus* sp.(Chlorophyta) extracted with different solvents." *Journal of Applied Phycology* 31 (2019): 1675-1683. <https://doi.org/10.1007/s10811-018-1726-5>
- [37] Piotrowska-Niczyporuk, Alicja, Andrzej Bajguz, Marta Talarek, Monika Bralska, and Elżbieta Zambrzycka. "The effect of lead on the growth, content of primary metabolites, and antioxidant response of green alga *Acutodesmus obliquus* (Chlorophyceae)." *Environmental Science and Pollution Research* 22 (2015): 19112-19123. <https://doi.org/10.1007/s11356-015-5118-y>
- [38] Almendinger, Martin, Franziska Saalfrank, Sascha Rohn, Elke Kurth, Monika Springer, and Daniel Pleissner. "Characterization of selected microalgae and cyanobacteria as sources of compounds with antioxidant capacity." *Algal Research* 53 (2021): 102168. <https://doi.org/10.1016/j.algal.2020.102168>
- [39] Amzah, Nur Faiqah, Shaza Eva Binti Mohamad, Fazrena Nadia Md Akhir, Fatin Syahirah Othman, Iwane Suzuki, and Yoshiaki Maeda. "Effect of agitation on growth and fatty acid composition of green microalgae *Acutodesmus obliquus* Q2-12E." *Chemical Engineering Transactions* 106 (2023): 1231-1236.
- [40] Aremu, Adeyemi O., Nqobile A. Masondo, Zoltan Molnár, Wendy A. Stirk, Vince Ördög, and Johannes Van Staden. "Changes in phytochemical content and pharmacological activities of three *Chlorella* strains grown in different nitrogen conditions." *Journal of applied phycology* 28 (2016): 149-159. <https://doi.org/10.1007/s10811-015-0568-7>
- [41] Custódio, Luísa, Fernando Soares, Hugo Pereira, Luísa Barreira, Catarina Vizetto-Duarte, Maria João Rodrigues, Amélia Pilar Rauter, Fernando Alberício, and João Varela. "Fatty acid composition and biological activities of *Isochrysis galbana* T-ISO, *Tetraselmis* sp. and *Scenedesmus* sp.: Possible application in the pharmaceutical and functional food industries." *Journal of applied phycology* 26 (2014): 151-161. <https://doi.org/10.1007/s10811-013-0098-0>
- [42] Talero, Elena, Sofía García-Mauriño, Javier Ávila-Román, Azahara Rodríguez-Luna, Antonio Alcaide, and Virginia Motilva. "Bioactive compounds isolated from microalgae in chronic inflammation and cancer." *Marine drugs* 13, no. 10 (2015): 6152-6209. <https://doi.org/10.3390/md13106152>
- [43] Elia, Josiane, Karina Petit, Jean-Michel Huvelin, Mona Tannoury, Mona Diab-Assaf, Delphine Carbonnelle, and Hassan Nazih. "Acetone fraction of the red marine alga *Laurencia papillosa* reduces the expression of Bcl-2 Anti-apoptotic marker and flotillin-2 lipid raft marker in MCF-7 breast cancer cells." *Iranian Journal of Pharmaceutical Research: IJPR* 19, no. 1 (2020): 321.
- [44] Goh, Su Hua, Noorjahan Banu Mohamed Alitheen, Fatimah Md Yusoff, Swee Keong Yap, and Su Peng Loh. "Crude ethyl acetate extract of marine microalga, *Chaetoceros calcitrans*, induces Apoptosis in MDA-MB-231 breast cancer cells." *Pharmacognosy Magazine* 10, no. 37 (2014): 1. <https://doi.org/10.4103/0973-1296.126650>

- [45] Suh, Sung-Suk, Eun Jin Yang, Sung Gu Lee, Ui Joung Youn, Se Jong Han, Il-Chan Kim, and Sanghee Kim. "Bioactivities of ethanol extract from the Antarctic freshwater microalga, *Chloromonas* sp." *International journal of medical sciences* 14, no. 6 (2017): 560. <https://doi.org/10.7150/ijms.18702>
- [46] Ebrahimi Nigjeh, Siyamak, Fatimah Md Yusoff, Noorjahan Banu Mohamed Alitheen, Mehdi Rasoli, Yeap Swee Keong, and Abdul Rahman bin Omar. "Cytotoxic effect of ethanol extract of microalga, *Chaetoceros calcitrans*, and its mechanisms in inducing apoptosis in human breast cancer cell line." *BioMed research international* 2013, no. 1 (2013): 783690. <https://doi.org/10.1155/2013/783690>
- [47] Murphy, Catherine, Sarah Hotchkiss, Jenny Worthington, and Stephanie R. McKeown. "The potential of seaweed as a source of drugs for use in cancer chemotherapy." *Journal of Applied Phycology* 26 (2014): 2211-2264. <https://doi.org/10.1007/s10811-014-0245-2>
- [48] Harizani, Maria, Efsthia Ioannou, and Vassilios Roussis. "The Laurencia paradox: An endless source of chemodiversity." *Progress in the chemistry of organic natural products* 102 (2016): 91-252. https://doi.org/10.1007/978-3-319-33172-0_2
- [49] Li-Beisson, Yonghua, Jay J. Thelen, Eric Fedosejevs, and John L. Harwood. "The lipid biochemistry of eukaryotic algae." *Progress in Lipid Research* 74 (2019): 31-68. <https://doi.org/10.1016/j.plipres.2019.01.003>
- [50] Niazi, Shaik Kalimulla, Dhanyakumara Shivapoojar Basavarajappa, Sushma Hatti Kumaraswamy, Asmatanzeem Bepari, Halaswamy Hiremath, Shashiraj Kariyellappa Nagaraja, Muthuraj Rudrappa, Anil Hugar, Mary Anne Wong Cordero, and Sreenivasa Nayaka. "GC-MS based characterization, antibacterial, antifungal and anti-oncogenic activity of ethyl acetate extract of *Aspergillus niger* strain AK-6 isolated from rhizospheric soil." *Current Issues in Molecular Biology* 45, no. 5 (2023): 3733-3756. <https://doi.org/10.3390/cimb45050241>
- [51] Swantara, Made Dira, Wiwik Susanah Rita, Nyoman Suartha, and Kadek Karang Agustina. "Anticancer activities of toxic isolate of *Xestospongia testudinaria* sponge." *Veterinary World* 12, no. 9 (2019): 1434. <https://doi.org/10.14202/vetworld.2019.1434-1440>
- [52] Renukadevi, K. P., P. S. Saravana, and J. Angayarkanni. "Antimicrobial and antioxidant activity of *Chlamydomonas reinhardtii* sp." *International journal of Pharmaceutical Sciences and Research* 2, no. 6 (2011): 1467.
- [53] Berneira, Lucas M., Ivandra I. de Santi, Caroline C. da Silva, Dalila Venzke, Pio Colepicolo, Rodrigo de A. Vaucher, Marco AZ Dos Santos, and Claudio MP de Pereira. "Bioactivity and composition of lipophilic metabolites extracted from Antarctic macroalgae." *Brazilian Journal of Microbiology* 52 (2021): 1275-1285. <https://doi.org/10.1007/s42770-021-00475-6>
- [54] Ozdemir, Guven, N. Ulku Karabay, Meltem Conk Dalay, and Baris Pazarbasi. "Antibacterial activity of volatile component and various extracts of *Spirulina platensis*." *Phytotherapy Research: An International Journal devoted to pharmacological and toxicological evaluation of natural product derivatives* 18, no. 9 (2004): 754-757. <https://doi.org/10.1002/ptr.1541>
- [55] Radman, Sanja, Martina Čagalj, Vida Šimat, and Igor Jerković. "Seasonal monitoring of volatiles and antioxidant activity of brown alga *Cladostephus spongiosus*." *Marine drugs* 21, no. 7 (2023): 415. <https://doi.org/10.3390/md21070415>
- [56] Kumari, Nalinee, Ekta Menghani, and Rekha Mithal. "Bioactive compounds characterization and antibacterial potentials of actinomycetes isolated from rhizospheric soil." (2019).
- [57] Yadav, A., M. Yadav, S. Kumar, D. Sharma, and J. P. Yadav. "In vitro antioxidant activities and GC-MS analysis of different solvent extracts of *Acacia nilotica* leaves." *Indian J. Pharm. Sci* 80, no. 5 (2018): 892-902. <https://doi.org/10.4172/pharmaceutical-sciences.1000436>
- [58] Pacheco, Bruna Silveira, Marco Aurélio Ziemann Dos Santos, Eduarda Schultze, Rosiane Mastelari Martins, Rafael Guerra Lund, Fabiana Kömmling Seixas, Pio Colepicolo, Tiago Collares, Favero Reisdorfer Paula, and Claudio Martin Pereira De Pereira. "Cytotoxic activity of fatty acids from Antarctic macroalgae on the growth of human breast cancer cells." *Frontiers in bioengineering and biotechnology* 6 (2018): 185. <https://doi.org/10.3389/fbioe.2018.00185>
- [59] de Andrade Tomaz, Anna Cláudia, George Emmanuel Cavalcanti de Miranda, Maria de Fátima Vanderlei de Souza, and Emídio Vasconcelos Leitão da Cunha. "Analysis and characterization of methyl esters of fatty acids of some *Gracilaria* species." *Biochemical Systematics and Ecology* 44 (2012): 303-306. <https://doi.org/10.1016/j.bse.2012.02.006>
- [60] Soni, Khushaboo, Preeti Maurya, and Sanjay Singh. "Bioactive compounds from fresh water green macro algae of Ganga water." *Journal of Pharmacognosy and Phytochemistry* 12, no. 5 (2023): 369-385. <https://doi.org/10.22271/phyto.2023.v12.i5d.14750>
- [61] Parimalachelvam, G., N. Nagendran, and S. Balakrishnan. "Assessment of anti-cancer properties of *Chlorella sorokiniana* against A375, A549 and HeLa cell lines using GC-MS analysis and MTT assay." *Sci Academique* 4, no. 1 (2023): 55-71.
- [62] Bharath, Boobalan, Kantharaj Perinbam, Sandhanasamy Devanesan, Mohamad S. AlSalhi, and Muthupandian

- Saravanan. "Evaluation of the anticancer potential of Hexadecanoic acid from brown algae *Turbinaria ornata* on HT-29 colon cancer cells." *Journal of Molecular Structure* 1235 (2021): 130229. <https://doi.org/10.1016/j.molstruc.2021.130229>
- [63] Suresh, Arumugham, Ramasamy Praveenkumar, Ramasamy Thangaraj, Felix Lewis Oscar, Edachery Baldev, Dharumadurai Dhanasekaran, and Nooruddin Thajuddin. "Microalgal fatty acid methyl ester a new source of bioactive compounds with antimicrobial activity." *Asian Pacific Journal of Tropical Disease* 4 (2014): S979-S984. [https://doi.org/10.1016/S2222-1808\(14\)60769-6](https://doi.org/10.1016/S2222-1808(14)60769-6)
- [64] Chen, Weixian, Jiaming Wang, Yangyi Ren, Hui Chen, Chenliu He, and Qiang Wang. "Optimized production and enrichment of α -linolenic acid by *Scenedesmus* sp. HSJ296." *Algal Research* 60 (2021): 102505. <https://doi.org/10.1016/j.algal.2021.102505>
- [65] Vilakazi, Hlengiwe, Tosin A. Olasehinde, and Ademola O. Olaniran. "Chemical characterization, antiproliferative and antioxidant activities of polyunsaturated fatty acid-rich extracts from *Chlorella* sp. S14." *Molecules* 26, no. 14 (2021): 4109. <https://doi.org/10.3390/molecules26144109>
- [66] Wiggins, Ashleigh KA, Julie K. Mason, and Lilian U. Thompson. "Growth and gene expression differ over time in alpha-linolenic acid treated breast cancer cells." *Experimental cell research* 333, no. 1 (2015): 147-154. <https://doi.org/10.1016/j.yexcr.2015.02.020>
- [67] Mason-Ennis, Julie K., Lauren P. LeMay-Nedjelski, Ashleigh KA Wiggins, and Lilian U. Thompson. "Exploration of mechanisms of α -linolenic acid in reducing the growth of oestrogen receptor positive breast cancer cells (MCF-7)." *Journal of functional foods* 24 (2016): 513-519. <https://doi.org/10.1016/j.jff.2016.04.037>