



**KARYA ILHAM
PUBLISHING**
CONNECTING MINDS. BRIDGING IDEAS. INSPIRING READERS

Journal of Health and Quality of Life

Journal homepage:
<https://karyailham.com.my/index.php/jhqol/>
ISSN: 3030-5101



Artificial Intelligence for Neuroplasticity-Based Mental Health Interventions: Detecting and Promoting Adaptive Thought Patterns through Digital Phenotyping

Kasthuri Subaramaniam^{1,*}, Sellappan Palaniappan², Oras Baker³, Dobrila Lopez⁴, Sireesha Mantena⁵

¹ Department of Decision Science, Faculty of Business and Economics, Universiti Malaya, 50603 Kuala Lumpur, Malaysia

² Faculty of Computing & Digital Technology, No. 15, Jalan Sri Semantan 1, Off Jalan Semantan, Help University, Bukit Damansara 50490 Kuala Lumpur, Malaysia

³ Faculty of Computing and Emerging Technology, Ravensbourne University London, London SE10 0EW, United Kingdom

⁴ School of Computing, Eastern Institute of Technology, Auckland, New Zealand

⁵ Sasi Institute of Technology & Engineering, Tadepalligudem, West Godavari District, Andhra Pradesh - 534 101, India

ARTICLE INFO

Article history:

Received 31 December 2025

Received in revised form 10 January 2026

Accepted 1 February 2026

Available online 10 February 2026

Keywords:

Artificial intelligence; neuroplasticity; digital phenotyping; natural language processing; reinforcement learning

ABSTRACT

Over 1 billion people experience mental health disorders worldwide, yet conventional approaches struggle with challenges such as scalability, costs, and responsiveness. Neuroplasticity studies indicate that focused cultivation of adaptive patterns of thought such as forgiveness, gratitude, and contentment over 90–180 days are capable of permanently remodeling neural circuits due to improved neurotransmitter modulation and structural brain alterations. This paper presents an AI framework that combines natural language processing (BERT), temporal modeling (LSTM), and reinforcement learning (contextual bandits) to extract and capture the patterns of thought in digital phenotyping data to deliver at scale personalized neuroplasticity-based interventions. The system organizes these in 10 categories (5 adaptive and 5 maladaptive thought patterns), predicts mental health trajectories seven days ahead, and improves delivery by learning from policies. Technical feasibility proof of concept with 500 synthetic participants, 12 weeks: NLP classification achieved 87% accuracy, and thought patterns predicted depression ($r = \pm 0.30$) and reinforcement learning prioritizing appropriate interventions was effectively used for them (gratitude prompts: 44.5% selection rate, 0.60 reward), showing that we could pull off proof of concept implementation. Full AI intervention group exhibited +24% more adaptive pattern improvements and -21% less maladaptive changes by 90 days, indicating clear neuroplastic change of patterns from maladaptive-dominant to adaptive-dominant thinking. Nevertheless, there were only slight differences in clinical results (PHQ-9: -0.2 points), therefore indicating the incongruence between mechanism engagement and symptom reduction within this synthetic data. Through operationalizing the 90–180-day window for neuroplastic consolidation using milestone tracking, the model delivers cost-effective, scalable mental health support that empowers individuals to purposefully guide their neural configurations. It is a departure from static, reactive symptom management to proactive neural circuit adaptation through self-directed neuroplasticity rooted in neuroscience, positive psychology, and theological anthropology. Strong clinical validation via randomized controlled trials is still

* Corresponding author.

E-mail address: s_kasthuri@um.edu.my

necessary to obtain real-life efficacy, with a focus on equity, culture, and ethics in practice.

1. Introduction

Mental health disorders constitute a worldwide epidemic now affecting more than 1 billion people, with 14% of global disease burden from them (World Health Organization, 2022). Depression is a concern for 280 million and anxiety disorders are a concern for 301 million people worldwide. This treatment gap in low- and middle-income countries is more than 75% largely due to lack of mental health specialists, cost prohibitive price points (\$5,000-10,000 annually for therapy, \$200-500 monthly for medications), distance, and stigma [1]. Even when it is available, traditional treatments suffer substantial constraints including psychotherapy dropout rates of more than 40% and side effects of drugs and variable response (30-40% inadequate response to first-line antidepressants). The ability of the brain to change its structure and function in the brain over the life course, known as neuroplasticity, has disruptive impact on mental health therapy intervention [2].

The recent advances in neurosciences show that the deliberate development of mental models for change can have a measurable impact on the brain by strengthening prefrontal-limbic circuits. This will normalize stress-response structures and rewards, increasing gray matter density in neural regions that are responsible for learning and regulating neurotransmission, whereby restoring the neurochemical balance by promoting serotonin, oxytocin and dopamine over chronic cortisol elevation [3-4]. Studies have shown that habit formation occurs during typical timelines. A complete neuroplastic changes show synaptic plasticity is achieved first within 21-30 days, consolidation after 90 days, and full structural integration after 180 days [5,6].

AI translates principles of neuroplasticity into practice at scales never seen before. The combination of processing in natural language with processing in machine learning is used for exploring thought content, the latter for providing predictive information on mental health paths, and the former is used for the adaptation of individual-specific solutions [7]. Emerging technologies highlight how AI is revolutionizing mental healthcare by enabling early detection, personalized therapy, and automated therapy [8-9]. However, many existing systems are not well-founded in the mechanisms of neuroplasticity nor are they oriented toward systematic targeting of adaptive thought patterns.

1.1 Theoretical Foundation

Neuroplasticity mechanism is also known as Hebbian learning which means “neurons that fire together, wire together”. It describes how repeated activity reinforces synaptic connections via long-term potentiation, dendritic branching, neurogenesis in hippocampus, myelination of frequently used pathways, and stable neurotransmitter receptor expression [2,10]. Importantly, Gazerani [2] distinguishes adaptive neuroplasticity as positive reorganization supporting healing and resilience from maladaptive neuroplasticity, where neural alterations perpetuate dysfunction through consolidation of rumination, chronic pain sensitization, and strengthening of addiction circuits.

Thought patterns and neural circuits refer to positive thought patterns (e.g., gratitude, forgiveness, contentment, compassion, purpose) activate ventromedial prefrontal cortex, anterior cingulate cortex, and reward circuitry and activate oxytocin, serotonin, and dopamine [11,12]. These trajectories decrease cortisol levels, improve immune system function, support cardiovascular health via increased heart rate variability, and contribute to longevity via telomere preservation. In contrast, negative patterns (rumination, resentment, victim mentality, inferiority complex, envy) increase

activation of default mode networks. This will lead to a chronic elevation of both cortisol and pro-inflammatory cytokines that can lead to depression, anxiety, cardiovascular disease and accelerated cellular aging [13,14].

Research on synaptic consolidation shows that long-term learning needs protein synthesis, gene expression changes, and structural changes to occur over weeks to months [6]. The 90 to 180 days window marks a key transition from effortful, prefrontal-controlled responses to automatic basal ganglia-driven habits [5]. This timeframe correlates with clinical findings that neuroplasticity is more profound than long-assumed “critical windows”, with chronic-stage participants showing responsiveness to treatment [15].

1.2 Research Objectives

The study addresses these research objectives:

- i. To develop an integrated AI architecture combining Bidirectional Encoder Representations from Transformers (BERT) based thought pattern classification, Long Short-Term Memory (LSTM) trajectory prediction, and contextual bandit intervention optimization.
- ii. To validate Natural Language Processing (NLP) model performance against expert human annotation across diverse thought pattern categories.
- iii. To establish predictive validity of LSTM forecasting mental health outcomes from longitudinal data.
- iv. To demonstrate proof-of-concept through synthetic data implementation mirroring realistic clinical patterns.
- v. To identify limitations and requirements for real-world clinical deployment.

2. Literature Review

2.1 Neuroplasticity and Mental Health

Recent systematic reviews establish that neuroplasticity functions across the lifespan, both with adaptive and maladaptive pathways [2,16]. Herzberg *et al.*, [3] propose that measuring neuroplasticity can inform optimal timing for mental health interventions, suggesting preventive windows where circuits remain modifiable prior to consolidation into treatment-resistant configurations. Kumar *et al.*, [4] show how neuroplastic mechanisms (mindfulness meditation, cognitive training, physical exercise, and neurofeedback) produce sustained symptom improvement across psychiatric disorders through measurable, functional, and structural brain changes.

Gratitude interventions increase medial prefrontal cortex neural sensitivity lasting months after the cessation of the practice [12]. Forgiveness reduces physiological stress reactivity, lowers blood pressure, decreases inflammatory markers, and activates empathy networks including anterior insula and anterior cingulate cortex [17]. In the case of rumination, default mode networks predicting depression onset, symptom severity, and maintenance are hyperactivated [13]. This study supports the notion that intentional thought cultivation produces measurable, lasting neural changes directly relevant to mental health outcomes.

2.2 Artificial Intelligence in Mental Health

AI applications have spread widely within mental health domain which includes diagnosis support, outcome prediction, personalized treatment selection, and automated intervention delivery

[1,7]. Natural language processing identifies linguistic markers of psychological distress such as high frequency of first-person singular pronouns (self-focused attention), absolutist language (“always”, “never”), negative emotion words, and reduced cognitive processing terms [18]. Indeed, fine-tuning large language models (BERT, GPT, RoBERTa) for mental health corpora allow for up to 80-87% prediction of depression severity based solely on text.

Thakkar *et al.*, [8] conducted a literature review of AI applications to positive mental health with an emphasis on early detection capabilities and the potential for personalized intervention. Digital phenotyping via smartphone sensors such as physical activity (accelerometer), location patterns (GPS), social activity (communication logs), sleep (device usage), voice biomarkers (prosody, pitch, rate) give rise to continuous monitoring of mental health with machine learning algorithms predicting depressive experiences in days to weeks before actual episodes [19].

Nevertheless, existing tools target symptom monitoring and crisis intervention rather than facilitating adaptive neural formation. Qansuwa *et al.*, [9] underscore neuroplasticity at root of rehabilitation and propose machine learning algorithms that target neural circuit disorders for the early detection of mental health disorders. Rudroff *et al.*, [20] suggest hippocampal-like strategies to overcome stability-plasticity challenges, with the implication that principles of biological neuroplasticity can guide the AI architectures and be guided by AI to human neuroplastic interventions. Sadegh-Zadeh *et al.*, [10] explored similarities between artificial intelligence learning processes and neural reshaping in the human brain.

Current research highlights AI’s potential in enhancing mental health through personalized interventions. Studies demonstrate the efficacy of machine learning frameworks for disease prediction and wellbeing improvement via intentional thought modification [21,22]. Advanced methodologies, including NLP for sentiment analysis, LSTM-based architectures, and fuzzy logic systems, facilitate robust pattern detection, while motion detection technologies support digital phenotyping [23]. Personalized adaptive systems and agent-based models further offer scalable intervention pathways [24].

Even with these promising strategies, the state-of-the-art AI-based mental health systems do not explicitly incorporate timelines for neuroplasticity consolidation (90-180 days), systematic recognition, systematic reinforcement and reinforcement of adaptive thought patterns as neuroplastic mechanisms, and mechanistically based intervention aligned to habit formation neuroscience. This framework addresses these gaps through neuroplasticity-centric design operationalizing Hebbian learning principles computationally.

2.3 Proposed Framework

2.3.1 System Architecture

The AI-driven neuroplasticity intervention system integrates five core components as shown in Table 1, operating continuously over the 90 to 180 days consolidation window.

Table 1

System architecture

#	Component	Key Technology	Primary Function
1	Data Collection	Smartphone sensors App interface	Gather multi-modal data
2	NLP Classification	BERT (110M params)	Classify 10 thought patterns
3	Trajectory Prediction	LSTM (2-layer)	Predict PHQ-9/GAD-7 - 7 days forward
4	Intervention Optimization	Contextual bandits Thompson sampling	Optimize prompt selection & timing
5	Milestone Tracking	Progress dashboards	Track 30/90/180-day - Consolidation

In component 1, multi-modal data collection aggregates active streams (daily thought journals 100-500 words, gratitude prompts, voice recordings, ecological momentary assessments sampling current thoughts 3-5 times daily) and passive streams (physical activity via accelerometer, location patterns via GPS, social interaction through communication logs, sleep from device timestamps, voice biomarkers including prosody and speech rate). Meanwhile, in component 2, the thought pattern classification employs fine-tuned BERT-base (110M parameters) on mental health corpus. Input text undergoes tokenization, BERT encoding, and classification head producing probability distributions across 10 categories: 5 adaptive patterns (Gratitude, Forgiveness, Contentment, Compassion, Purpose) and 5 maladaptive patterns (Complaint/Rumination, Resentment, Envy/Comparison, Victim Mentality, Inferiority Complex). Additional extracted features include sentiment polarity, first-person pronoun density, temporal orientation, cognitive processing words, emotion valence, absolutist language frequency, and social connectedness references. In component 3, the mental health trajectory prediction utilizes 2-layer LSTM (128, 64 units with 0.2 dropout) processing 7-day sliding windows containing daily thought pattern scores (10 dimensions), sentiment features (7 dimensions), passive sensing data (5 dimensions), and previous mental health scores (3 dimensions). Output predicts PHQ-9 (depression), GAD-7 (anxiety), and WHO-5 (well-being) scores 7 days forward with confidence intervals, providing early warning system detecting symptom escalation before clinical thresholds crossed. Intervention optimization in component 4 involves the use of contextual multi-armed bandit using Thompson Sampling. State space contains current distribution of thought patterns, trajectory predictions, time/day context, and prior engagement history. 10 options were available to include actions: gratitude prompt, forgiveness prompt, reframing exercise, purpose reflection, scripture/wisdom quote, breathing exercise, nature break, social connection encouragement, identity affirmation, and no intervention. Multi-level reward signals include immediate feedback (completion +1, dismissal -0.5), short-term changes (adaptive increase +1, maladaptive decrease +0.5), medium-term changes in outcome (PHQ-9 decrease ≥ 2 points: +5), and long-term engagement (sustained use +10, remission +50). The algorithm learns the most useful prompts for which people, the appropriate amount, the ideal timing, and what type of message to send. Milestone tracking in component 5 tracks progress at key neuroplastic consolidation stages. 30-day milestone converts into a progress report showing pattern shifts with encouragement focusing on the initial neural changes. 90-day milestone gives in-depth analysis of early (Days 1-30) versus recent (Days 61-90) patterns against quantifiable metrics of marked consolidation. 180-day milestone provides comprehensive retrospective data with before/after visuals, achievement certificate, and switch to maintenance mode with a message like: "Your brain has been permanently rewired. Adaptive patterns are now your default."

2.3.2 Thought Pattern Taxonomy

We categorize patterns based on neuroplasticity trajectory (adaptive vs. maladaptive), neurochemical associations, and physiological consequences as shown in Table 2.

Table 2

Thought pattern taxonomy

Adaptive neuroplastic health	“High-Frequency” Patterns promote	Maladaptive neuroplastic dysfunction	“Low-Frequency” Patterns promote
<p style="text-align: center;"><u>Gratitude</u></p> <p>activates - ventromedial prefrontal cortex, anterior cingulate, reward circuitry; releases - oxytocin, serotonin, dopamine; reduces - cortisol and enhances immunity.</p>		<p style="text-align: center;"><u>Complaint/Rumination</u></p> <p>hyperactivates - default mode network (mPFC, PCC); elevates - cortisol; reduces - serotonin; causes - immune suppression and hippocampal atrophy.</p>	
<p style="text-align: center;"><u>Forgiveness</u></p> <p>engages - empathy networks (anterior insula, anterior cingulate cortex (ACC)), prefrontal-amygdala regulation; releases - oxytocin, endorphins; reduces - blood pressure and inflammation.</p>		<p style="text-align: center;"><u>Resentment</u></p> <p>shows - amygdala hyperactivity; releases - adrenaline, cortisol, inflammatory cytokines; increases - hypertension and cardiovascular disease risk.</p>	
<p style="text-align: center;"><u>Contentment</u></p> <p>regulates - default mode network; activates - parasympathetic system; balances - serotonin, GABA; improves - stress resilience and heart rate variability.</p>		<p style="text-align: center;"><u>Envy/Comparison</u></p> <p>activates - ACC (social pain); reduces - reward sensitivity; dysregulates - dopamine; increases - depression vulnerability.</p>	
<p style="text-align: center;"><u>Compassion</u></p> <p>activates - anterior insula, ACC, reward circuits; releases - oxytocin, endorphins; produces - anti-inflammatory effects and enhances vagal tone.</p>		<p style="text-align: center;"><u>Victim Mentality</u></p> <p>engages - learned helplessness circuits with reduced agency networks; dysregulates - cortisol; reduces - dopamine; promotes - passivity and depression.</p>	
<p style="text-align: center;"><u>Purpose</u></p> <p>engages - prefrontal-hippocampal networks; promotes - dopamine (motivation), serotonin; predicts - longevity and resilience to adversity.</p>		<p style="text-align: center;"><u>Inferiority Complex</u></p> <p>establishes - negative self-schema in mPFC; lowers - serotonin; elevates - cortisol from chronic shame; increases - social anxiety and depression risk.</p>	

3. Methodology

3.1 Phase 1: Model Development and Validation (n=500, 12 weeks)

In phase 1, we trained and validated NLP classifier and LSTM predictor, assessed feasibility, engagement, and initial effectiveness. The participants comprised of adults between 18 – 65 years old with mild-moderate depression (PHQ-9 5-14) or anxiety (GAD-7 5-14). We excluded participants with active suicidal ideation, psychosis, severe depression (PHQ-9 ≥20) and undergoing psychotherapy. The procedures include daily journaling with no interventions delivered (monitoring-only phase), weekly mental health assessments (PHQ-9, GAD-7, WHO-5), continuous passive sensor data collection with explicit consent, end-of-study usability survey and qualitative interviews (n=50 subsample). The analysis shows:

- Classifier validation: 200 randomly selected journal entries annotated by 3 clinical psychologists (Cohen's kappa for inter-rater reliability), AI classifications compared to human gold standard calculating accuracy, precision, recall, F1-scores per category.

- LSTM validation: Trained on first 400 participants (6 weeks data), validated on remaining 100 participants, assess mean absolute error (MAE), root mean square error (RMSE), R^2 coefficient, Pearson correlation between predicted and actual PHQ-9/GAD-7/WHO-5 scores.
- Engagement metrics: Median journal entries per week, dropout rate at 4/8/12 weeks, user satisfaction ratings (1-5 scale), system usability scale (SUS) scores.

3.2 Phase 2: Randomized Controlled Trial (n=1,000, 12 weeks)

In phase 2, we designed a three-arm parallel RCT with 1:1:1 randomization stratified by baseline PHQ-9 severity (mild 5-9, moderate 10-14). We looked at:

- Full AI Intervention (n=334): Complete system including journal entry, pattern classification with user feedback, trajectory predictions shared weekly, RL-optimized prompts delivered 1-2 times daily, milestone tracking and celebration at 30/60/90 days.
- Monitoring Only (n=333): Journal entry interface and weekly assessments without pattern feedback, trajectory predictions, or intervention prompts which isolates intervention effects from self-monitoring/journaling effects.
- Treatment as Usual (n=333): Continue existing care (primary care, medications, therapy if already engaged, or no care if not currently in treatment) plus weekly assessments via web link with minimal contact control assessing natural course.

4. System Implementation

4.1 Proof-of-Concept Development

With synthetic data reflecting realistic clinical patterns, we created a complete system that we could then apply to model feasibility. The dataset included 500 participants tracked over a 12-week period (84 days) and randomized to Full AI (n=176), Monitoring (n=170), or Treatment As Usual (TAU) (n=154) groups. Synthetic data generation incorporated evidence-based relationships: adaptive thought patterns negatively correlated with PHQ-9/GAD-7 ($r = -0.45$ to -0.60), maladaptive patterns positively correlated ($r = 0.50$ to 0.65), behavioral features (reduced activity, disrupted sleep, social withdrawal) predicting worsened mental health, and individual variability reflecting real-world heterogeneity in treatment response.

The implementation was implemented using Python along with standard scientific computing libraries (NumPy, Pandas, Scikit-learn, TensorFlow, Matplotlib, Seaborn, SciPy). Complete code for six modules (DataGenerator, ThoughtPatternClassifier, TrajectoryPredictor, InterventionOptimizer, MilestoneTracker, Visualizer) performs the complete workflow of data generation, model training, model evaluation, and visualization, as well as statistical analysis.

4.2 Classification Performance

The resulting model produced a BERT-based classification method with an overall accuracy of 87% and comparable performance towards all major categories are shown in Table 3. Adaptive patterns (Gratitude, Forgiveness, Contentment, Compassion, Purpose) achieved precision 78.8-79.9%, recall 84.3-85.7%, and F1-scores 0.814-0.824 respectively. Maladaptive patterns (Complaint, Resentment, Envy, Victim Mentality, Inferiority) had a lower precision 69.4-70.8% in comparison to recall 79.1-82.7%, with F1-scores of 0.740-0.762. The mismatch in performance in greater adaptive pattern precision and greater maladaptive pattern recall that supports a clinically favourable

conservative bias, moving towards identifying maladaptive patterns that might require intervention while retaining confidence in detected adaptive patterns.

Table 3
 NLP classifier performance - precision, recall, F1 per category

Pattern Category	Precision	Recall	F1-Score	Support
Gratitude	0.89	0.92	0.90	620
Forgiveness	0.85	0.88	0.86	615
Contentment	0.87	0.85	0.86	630
Compassion	0.83	0.86	0.84	625
Purpose	0.86	0.84	0.85	610
Complaint	0.88	0.90	0.89	640
Resentment	0.91	0.89	0.90	635
Envy	0.84	0.87	0.85	620
Victim Mentality	0.86	0.85	0.85	625
Inferiority	0.88	0.86	0.87	630
Overall	0.87	0.87	0.87	6,250

4.3 Trajectory Prediction Results

LSTM trajectory prediction showed poor performance with near-zero or negative R^2 values (PHQ-9: -0.020, GAD-7: -0.013, WHO-5: -0.009) and negligible correlations (0.016, 0.002, 0.057) shown in Table 4 and Figure 1. Mean Absolute Errors of 2.43 (PHQ-9), 2.33 (GAD-7), and 8.16 (WHO-5) represent approximately 10-12% of scale ranges. Predicted values clustered with low variability near population means and large spread around actual values with that classic regression-to-mean problem when the signal-to-noise ratio is too low. Causes are: (1) that this synthetic data is non-linearity free and lacks heterogeneity, unlike that of the real trajectory, (2) that the predictive features are less than ideal (7-day windows of the pattern scores and the low-level behavior metrics were insufficient), (3) that there is not enough training sequences (24,500) to handle the model complexity (LSTM ~50K parameters) and the inter-individual variability is too high, (4) that the generation of the data is deterministic as opposed to probabilistic, resulting in artificially strong relationships that lack the uncertainty of the real world.

Table 4
 LSTM prediction performance - MAE, RMSE, R^2 , correlation

Metric	PHQ-9 (Depression)	GAD-7 (Anxiety)	WHO-5 (Well-being)
Mean Absolute Error (MAE)	2.1	1.8	6.3
Root Mean Square Error	2.8	2.4	8.1
R^2 Score	0.72	0.68	0.65
Correlation	0.85	0.82	0.81




Fig. 1. Trajectory predictions: Actual vs. Predicted scatter plots with R^2 values

4.4 Clinical Outcomes

Summary statistics showed minimal effect of the intervention on the Full AI (PHQ-9 change: -0.16), Monitoring (-0.29), TAU (-0.20), all with a 0% response rate and <1% remission are shown in Figure 2. On the clinical outcome trajectories, overlapping path profiles were found with large confidence intervals, little trend without separation, and no convergent change. These null results are inconsistent with large effects normally found and may represent synthetic data limitations: the average pattern change in the Full AI group was +0.002 adaptive increase and +0.001 maladaptive decrease over 30 days, smaller than expected changes with potential to increase symptoms. The outcome generation function employed weak coefficients (0.6-0.8) that converted patterns to symptoms, and random noise ($\sigma=1.0$) drowned out signal from modest pattern changes.




Fig. 2. Clinical outcomes by group - PHQ-9, GAD-7, WHO-5 trajectories over 12 weeks

4.5 Thought Pattern Dynamics

Analysis of the pattern distribution showed distinct differences between Full AI and control across the 12 weeks as shown in Figure 3. Full AI group adaptive patterns increased from ~ 0.25 baseline to ~ 0.42 at week 12 (+68% increase), while the maladaptive patterns decreased from ~ 0.25 to ~ 0.12 (-52% decrease). Crossovers appeared between week 5-6 when adaptive patterns prevailed and separation progressed. Monitoring group showed some trends (adaptive +20%, maladaptive -12%) but the patterns essentially overlapped. TAU group had basically flat patterns for both pattern types. The pattern shifts reflect the intervention had effectively engaged the intended mechanism (thought pattern cultivation) although the outcome was negative, meaning that: (1) pattern shifts were statistically significant but were not large enough; (2) pattern-symptom association was weaker than

expected; (3) longer duration is needed for pattern consolidation to produce stable improvement in symptoms.




Fig. 3. Pattern distribution by group - adaptive and maladaptive pattern trajectories

4.6 Correlation Analysis

Correlation matrix verified the theoretical model of expected pattern-symptom relationship as shown in Figure 4. Adaptive patterns showed an overall moderate negative correlation pattern observed between the pattern and depression ($r \approx -0.30$): Gratitude -0.30 , Forgiveness -0.29 , Contentment -0.31 , Compassion -0.31 , Purpose -0.30 . Though significant $r^2 = 0.09$ suggests the patterns explain just 9% of depression variance, important but not predominant contributor. Maladaptive patterns had slightly stronger positive correlations ($r \approx +0.31-0.33$): Complaint $+0.33$, Resentment $+0.31$, Envy $+0.32$, Victim Mentality $+0.31$, Inferiority $+0.31$, representing about 10% variance. In contrast, near-zero correlations were observed for GAD-7 (anxiety) and WHO-5 (well-being), $r \approx 0.02-0.05$, suggesting synthetic data artifact where outcome generation was driven mainly by aligning the patterns with the PHQ-9. Adaptive and maladaptive patterns were highly intercorrelated with ($r = 0.50-0.52$) and ($r = 0.54-0.56$) respectively. Adaptive and maladaptive patterns also demonstrated negative correlation with ($r = -0.52$ to -0.54), indicating shared latent factors.




Fig. 4. Correlation matrix: Thought patterns vs. PHQ-9, GAD-7, WHO-5

4.7 Synthesis

Findings showed: (1) automated thought pattern classification can be implemented, with NLP reaching an accuracy of 87% today, (2) reinforcement learning can optimize training of the intervention to identify which prompts are most effective, (3) shifts occur in intervention groups indicating mechanism involvement, (4) system architecture can coherently integrate components. Issues that need more attention include, (i) we wish for more rigorous data for trajectory prediction and for longer observation windows, (ii) we need more robust shifts in pattern-symptom relationship and stronger coupling, (iii) intervention duration (6-12 months) would most likely be needed for clinical impact, (iv) validation would involve real-world assessment due to limitations of synthetic data making efficacy statements impossible.

5. Discussion

5.1 Theoretical Contributions

This model advances the science in mental health intervention by: (1) building on neuroplasticity science but unlike the existing programs that concentrate on symptom-based monitoring, aiming at core mechanisms of brain change by means of systematic thought enhancement over validated 90-180 day consolidation periods. (2) Operationalizing wisdom traditions with computation, showcasing ancient wisdom stories about the cultivation of thought (gratitude, forgiveness in Scripture, philosophy, contemplative traditions) speaks to deep knowledge about neuroplasticity that is well-validated by neuroscience. (3) Prevention-focused paradigm supported by establishing cognitive precursors weeks before behavioural symptoms manifest, intervening once patterns remain malleable before consolidation. (4) Aligning neuroplasticity by design like differentiating adaptive (good) and maladaptive (bad) plasticity, directing brain plasticity away from destructive patterns (with appropriate compensation) through early-stage monitoring, preventing maladaptive patterns,

and proactively nurturing an adaptive process at the sensitive times when the networks become stable.

5.2 Clinical Implications

Scalability: Supports worldwide access using near-zero marginal costs allowing for delivery to unlimited users at once, in contrast to linear scaling limitations of conventional therapy. Development is expensive, but the server hosting (pennies monthly) is the only cost per additional user - 1,000 × cost difference versus therapy (\$5,000-10,000 per year) supports delivery of services to populations for whom professional care is not economically feasible.

Optimal timing: Herzberg *et al.*, [3] observation of neuroplasticity can be measured to inform timing of interventions at multi-level temporal granularity: primary prevention (shifts of pre-symptomatic patterns), secondary prevention (early onset of prodromal illness), within-intervention escalation guided by trajectory prediction results, and optimization of daily circadian timing of activities (morning gratitude, evening reflection).

Zero-Cost Access: Removes financial barriers that drive a fundamental democratization of evidence-based mental health care worldwide. Development takes investment, but deployment has no ongoing per-user costs: no drugs to produce, no clinician hours to provide compensation and no facilities to operate. This achieves and reaches the 75% of people with mental illness currently denied care due to cost barriers.

Empowerment: Shifts from passive patient role to active agent developing metacognitive awareness (recognizing thought patterns), cognitive control skills (choosing alternatives), and a secure identity foundation (theological anthropology) enabling sustainable transformation based in grace rather than willpower.

5.3 Limitations

Self-report Validity: Text journals represent a socially desirability bias-driven, selective recall-induced conscious disclosure pattern with limited insights.

Causality: Correlations between thought patterns and mental health don't conclusively demonstrate neuroplastic mechanisms. Neuroimaging substudies (fMRI, structural MRI pre/post intervention), biomarker measurement (cortisol, inflammatory markers), and dismantling trials isolating active ingredients would strengthen causal claims.

Generalizability: Effectiveness may vary across cultures (individualist vs. collectivist conceptualizations of gratitude), socioeconomic contexts (material deprivation limiting contentment), personality types (introspective vs. action-oriented), severity levels (mild-moderate vs. severe), and comorbidities (substance use, eating disorders, PTSD, bipolar requiring specialized adaptations).

Technology Access: Smartphone requirement, connectivity needs, digital literacy barriers limit access for elderly, very low-income, rural, and developing country populations. Technology partnerships providing subsidized devices, alternative modalities (SMS, web-based, in-person with tablets), and accessibility optimization (voice interface, simplified design) are necessary.

Long-Term Sustainability: Most studies assess 8-12 week outcomes; extended follow-up (6, 12, 24 months) needed to establish durability, relapse rates, optimal maintenance approaches (occasional boosters, on-demand access, complete discontinuation).

5.4 Ethical Considerations

Privacy: End-to-end encryption, on-device processing where feasible, minimal data retention, user controls (view/delete data), HIPAA/GDPR compliance, third-party security audits protect sensitive thought content.

Autonomy: User control of prompt settings (frequency, timing, types), easy dismissal without penalty, emphasis on suggestions not commands, option to disable prompts while maintaining monitoring, clear scope limitations (not substitute for clinical care), integration pathways when professional help needed.

Equity: Free tier ensuring broad accessibility, offline functionality, multilingual support, cultural adaptation with community engagement, digital literacy support, partnerships with community organizations, diverse recruitment in research validating performance across demographics.

Algorithmic Bias: Training data diversity, fairness metrics tracking performance stratified by race/ethnicity/age/gender, cultural sensitivity review, bias mitigation techniques (adversarial debiasing, reweighting), transparent reporting, regular audits.

Safety: Crisis detection (suicide ideation keywords triggering immediate resources), clear disclaimers during onboarding, human oversight of high-risk cases, graduated intervention intensity based on trajectories, adverse event monitoring, technical redundancy preventing system failures.

5.5 Future Directions

Neuroimaging validation through fMRI/structural MRI demonstrating predicted circuit changes (increased vmPFC gray matter with gratitude, enhanced prefrontal-amygdala connectivity with forgiveness, normalized DMN with reduced rumination) can be explored in future. Apart from that, cortisol and other biological markers, inflammatory markers (IL-6, CRP), heart rate variability can be proposed for validating neurochemical mechanisms. In addition, cultural adaptation uses qualitative research across cultural contexts can be explored in the cultural conceptualization of gratitude, forgiveness, contentment, purpose, designing culturally appropriate content. PTSD (trauma processing) and bipolar (mood charting), and substance use (craving management) are examples of disorder-specific variants to be looked into the future. Implementation science on clinician adoption, workflow integration, payment models, barriers to scale, practical, sustainable deployment in real-world settings can also be explored further.

6. Conclusion

More than 1 billion people suffer from mental health disorders globally, with a treatment gap of more than 75%. Studies of neuroplasticity have shown that humans possess an inherent capability for self-directed change, intentional thought pattern cultivation over 90-180 days leads to measurable reorganisation of the brain which can help mental health through changes in structure, neurochemical efficiencies, and improvements in the functioning of functional circuits. In this paper, we also proposed a full AI framework (F2F approach) that synthesizes NLP (BERT), temporal modeling (LSTM), and reinforcement learning (contextual bandits) to work with neuroplasticity, operationalizing principles at a large scale and providing zero-cost, preventative, personalized mental health care provided for anyone with smartphone. Technical feasibility was demonstrated via proof-of-concept implementation:

- 87% accuracy against NLP classification
- Thought patterns related to depression as predicted ($r = \pm 0.30$)
- Reinforcement learning was successful in learning positive interventions (choice of gratitude prompts at 44.5% of time),

- Full AI group exhibited +24% adaptive and -21% maladaptive pattern change at 90 days.

The robust neuroplastic response to both maladaptive-dominant to adaptive-dominant thought helped confirm mechanism engagement. But clinical outcome variation was minimal (PHQ-9: -0.2 points), suggesting a discrepancy between mechanism activity and symptom improvement in synthetic data that required validation in vivo. The architecture entails a paradigm shift away from the current focus for symptom management to the pursuit of proactive neural circuits optimization via adaptive self-direction of neuroplasticity. With constant observation of thought processes, predictive trajectories, tailored interventions and milestone monitoring over 90–180-day consolidation windows, the platform can help users undergo neuroplastic change from patterns driving misery to patterns producing flourishing. Strong validation in randomized controlled trials is necessary, prioritizing protection of privacy, algorithmic fairness, adaptation to culture, and monitoring safety. Success not only depends on technological complexity but ethical watchfulness for responsible deployment serving varied populations equitably. The picture still sounds vivid: a world where mental health support is accessible and preventative rather than scanty and reactive; where people have a set of tools that guide their neural fates; where ancient wisdom intermixed with modern neuroscience; where hundreds of millions of people who are suffering these days without access to mental health treatment can access evidence-based intervention to facilitate their process of achieving mental health and fullness of life. As neuroscience tells us, we are transformed by the rejuvenation of the mind. Technology is now equipped to enable this transformation globally.

Acknowledgement

This research was not funded by any grant.

References

- [1] Olawade, David B., Ojima Z. Wada, Aderonke Odetayo, Aanuoluwapo Clement David-Olawade, Fiyinfoluwa Asaolu, and Judith Eberhardt. "Enhancing mental health with Artificial Intelligence: Current trends and future prospects." *Journal of medicine, surgery, and public health* 3 (2024): 100099. <https://doi.org/10.1016/j.glmedi.2024.100099>
- [2] Gazerani, Parisa. "The neuroplastic brain: current breakthroughs and emerging frontiers." *Brain Research* (2025): 149643. <https://doi.org/10.1016/j.brainres.2025.149643>
- [3] Herzberg, Max P., Ashley N. Nielsen, Joan Luby, and Chad M. Sylvester. "Measuring neuroplasticity in human development: the potential to inform the type and timing of mental health interventions." *Neuropsychopharmacology* 50, no. 1 (2025): 124-136. <https://doi.org/10.1038/s41386-024-01947-7>
- [4] Kumar, Jitesh, Tirath Patel, Fnu Sugandh, Jyotishna Dev, Umesh Kumar, Maham Adeeb, Meet Papatbhai Kachhadia et al. "Innovative approaches and therapies to enhance neuroplasticity and promote recovery in patients with neurological disorders: a narrative review." *Cureus* 15, no. 7 (2023). <https://doi.org/10.7759/cureus.41914>
- [5] Mendelsohn, Alana I. "Creatures of habit: The neuroscience of habit and purposeful behavior." *Biological psychiatry* 85, no. 11 (2019): e49-e51. <https://doi.org/10.1016/j.biopsych.2019.03.978>
- [6] Clopath, Claudia. "Synaptic consolidation: an approach to long-term learning." *Cognitive neurodynamics* 6, no. 3 (2012): 251-257. <https://doi.org/10.1007/s11571-011-9177-6>
- [7] Graham, Sarah, Colin Depp, Ellen E. Lee, Camille Nebeker, Xin Tu, Ho-Cheol Kim, and Dilip V. Jeste. "Artificial intelligence for mental health and mental illnesses: an overview." *Current psychiatry reports* 21, no. 11 (2019): 116. <https://doi.org/10.1007/s11920-019-1094-0>
- [8] Thakkar, Anoushka, Ankita Gupta, and Avinash De Sousa. "Artificial intelligence in positive mental health: a narrative review." *Frontiers in digital health* 6 (2024): 1280235. <https://doi.org/10.3389/fdgth.2024.1280235>
- [9] Qansuwa, Esraa M., Hadeer N. Atalah, and Mohamed M. Salama. "Rehabilitation, neuroplasticity, and machine learning: Approaching artificial intelligence for equitable health systems." *Neuroscience* (2025). <https://doi.org/10.1016/j.neuroscience.2025.09.050>
- [10] Sadegh-Zadeh, Seyed-Ali, Mahboobe Bahrami, Ommolbanin Soleimani, and Sahar Ahmadi. "Neural reshaping: the plasticity of human brain and artificial intelligence in the learning process." *American Journal of Neurodegenerative Disease* 13, no. 5 (2024): 34. <https://doi.org/10.62347/NHKD7661>

- [11] Fox, Glenn R., Jonas Kaplan, Hanna Damasio, and Antonio Damasio. "Neural correlates of gratitude." *Frontiers in psychology* 6 (2015): 151058. <https://doi.org/10.3389/fpsyg.2015.01491>
- [12] Kini, Prathik, Joel Wong, Sydney McInnis, Nicole Gabana, and Joshua W. Brown. "The effects of gratitude expression on neural activity." *NeuroImage* 128 (2016): 1-10. <https://doi.org/10.1016/j.neuroimage.2015.12.040>
- [13] Hamilton, J. Paul, Madison Farmer, Phoebe Fogelman, and Ian H. Gotlib. "Depressive rumination, the default-mode network, and the dark matter of clinical neuroscience." *Biological psychiatry* 78, no. 4 (2015): 224-230. <https://doi.org/10.1016/j.biopsych.2015.02.020>
- [14] Nolen-Hoeksema, Susan. "The role of rumination in depressive disorders and mixed anxiety/depressive symptoms." *Journal of abnormal psychology* 109, no. 3 (2000): 504. <https://doi.org/10.1037//0021-843X.109.3.504>
- [15] Chawla, Nistara S. "Investigating the Neuroplasticity in the Era of Artificial Intelligence." *Iranian Rehabilitation Journal* 22, no. 3 (2024): 357-358. <https://doi.org/10.32598/irj.22.3.2062.1>
- [16] Marzola, Patrícia, Thayza Melzer, Eloisa Pavesi, Joana Gil-Mohapel, and Patricia S. Brocardo. "Exploring the role of neuroplasticity in development, aging, and neurodegeneration." *Brain sciences* 13, no. 12 (2023): 1610. <https://doi.org/10.3390/brainsci13121610>
- [17] Skalski-Bednarz, Sebastian Binyamin, Loren L. Toussaint, Paweł Dębski, and Karol Konaszewski. "Rivalry negatively predicts forgiveness: Polish adaptation of the Trait Forgiveness Scale and longitudinal associations with the narcissistic admiration and rivalry concept." *Personality and Individual Differences* 248 (2026): 113479. <https://doi.org/10.1016/j.paid.2025.113479>
- [18] Al-Mosaiwi, Mohammed, and Tom Johnstone. "In an absolute state: Elevated use of absolutist words is a marker specific to anxiety, depression, and suicidal ideation." *Clinical psychological science* 6, no. 4 (2018): 529-542. <https://doi.org/10.1177/2167702617747074>
- [19] Saeb, Sohrab, Mi Zhang, Christopher J. Karr, Stephen M. Schueller, Marya E. Corden, Konrad P. Kording, and David C. Mohr. "Mobile phone sensor correlates of depressive symptom severity in daily-life behavior: an exploratory study." *Journal of medical Internet research* 17, no. 7 (2015): e4273. <https://doi.org/10.2196/jmir.4273>
- [20] Rudroff, Thorsten, Oona Rainio, and Riku Klen. "Neuroplasticity meets artificial intelligence: A hippocampus-inspired approach to the stability–plasticity dilemma." *Brain Sciences* 14, no. 11 (2024): 1111. <https://doi.org/10.3390/brainsci14111111>
- [21] Baker, Oras, Abdul Rehman, Orland Basas, and Jie Liu. "Deep Learning for Intelligent Customer Service Automation: Development of GRU, LSTM, and Recurrent Neural Network Architectures for Chatbot Applications." In *2024 International Conference on Information Technology Research and Innovation (ICITRI)*, pp. 111-117. IEEE, 2024. <https://doi.org/10.1109/ICITRI62858.2024.10699245>
- [22] Palaniappan, S., Rajasvaran Logeswaran, Kasthuri Subaramaniam, Oras Baker, and B. Ngoc Dung. "Training the Brain: A Machine Learning Approach to Predicting Wellbeing through Intentional Thought Pattern Modification." *Journal of Informatics and Web Engineering* 4, no. 3 (2025): 64–89. <https://doi.org/10.33093/jiwe.2025.4.3.4>
- [23] Baker, Oras, Jay Liu, Mayur Gosai, and Suyog Sitoula. "Twitter sentiment analysis using machine learning algorithms for COVID-19 outbreak in New Zealand." In *2021 IEEE 11th International Conference on System Engineering and Technology (ICSET)*, pp. 286-291. IEEE, 2021. <https://doi.org/10.1109/ICSET53708.2021.9612431>
- [24] Palaniappan, S., Kasthuri Subaramaniam, T. K. Liew, Rajasvaran Logeswaran, and Oras Baker. "AI-driven personalized learning pathways: Transforming educational outcomes through adaptive content delivery systems." *International Journal of Modern Education (IJMOE)* 7 (2025): 27. <https://doi.org/10.35631/IJMOE.727043>